

MAB TREATMENT OF MILD-MODERATE COVID-19 in confirmed positive patients

Initial Screening Criteria

- Within 10 days from symptom-onset
 - Adult ≥ 18 years of age
- OR
- Pediatric (12-17 years of age and weighs ≥ 40 kgs /88lbs)

Patient must meet the criteria above **PLUS** 1 of the following risk factors for disease progression:

- ≥ 65 years of age**
- 12-64 years of age** AND have
 - Body mass index (BMI) ≥ 25 or BMI ≥ 85 th percentile for their age and gender based on CDC growth charts
 - Chronic kidney disease or
 - Diabetes or
 - Cardiovascular disease/hypertension (including congenital heart disease) or
 - Chronic lung disease (COPD, mod-severe asthma, interstitial lung disease, cystic fibrosis, pulmonary HTN) or
 - Immunosuppressive disease* or
 - Currently receiving immunosuppressive treatment (chemotherapy, transplant immunosuppressants, immune modulators such as Rituximab, etc.)
 - Sickle cell disease or
 - Neurodevelopmental disorders or other conditions that confer medical complexity or
 - Medical-related technological dependence (tracheostomy, gastrostomy, or PPV not related to COVID-19) or
 - Pregnant

*AIDS or CD4 count < 200 , Complement deficiency, Grafts-Vs-Host disease (GVHD), HIV infection, Immunoglobulin deficiency/ Immunodeficiency, Immunosuppressive therapy (within the last 12 months), Leukemia, Lymphoma (Hodgkin's/ Non-Hodgkin's (NHL)), Metastatic cancer, Multiple Myeloma, Solid organ malignancy, Steroid therapy (within past 2 weeks), Bone marrow transplant (BMT) or peripheral stem cell transplant (PSCT), Solid organ transplant